Non-local Models in Replicator Dynamics

Nikos Kavallaris
(Joint work with J. Lankeit, M. Winkler)

KAAS, Karlstads University
30-11-2022

Evolutionary Game Theory

Evolutionary game theory (EGT) is the application of game theory to evolving populations in biology. It de nes a framework of contests, strategies, and analytics into which Darwinian competition can be modelled. It originated in 1973 with John Maynard Smith and George R. Price's formalisation of contests analysed as strategies, and the mathematical criteria that can be used to predict the results of competing strategies (Smith & Price (1973)).
Evolutionary game theory differs from classical game theory in focusing more on the dynamics of strategy change

Replicator Dynamics(Key Ideas)

Finite Strategy Space

1. The dynamics most widely used and studied in the literature on evolutionary game theory is the replicator dynamics.
2.

Replicator Dynamics (Pay-off)

Replicator dynamics (relevant equations)

1. Consider symmetric games with m finitely many players then we can assume that $p_i(t)$ evolve as differentiable functions.
2. For biological populations the above assumption means that infinitely big populations are considered and their generations blend continuously to each other.
3. A reasonable assumption, which is also in agreement with the basic tenet of Darwinism, is that per capita rate of growth (i.e. the logarithmic derivative) $\dot{p}_i/p_i$ is given by the difference between the pay-off for strategy i and the average pay-off. This yields the replicator dynamics system,

$$\dot{p}_i := \frac{dp_i}{dt} = \left(\sum_{j=1}^{m} a_{ij} p_j(t) \right) - \sum_{i=1}^{m} \sum_{j=1}^{m} a_{ij} p_i(t) p_j(t) \, p_i(t); \quad i = 1, 2, \ldots, m; \quad t > 0; \qquad (1)$$

   (Bomze (1990)).
4. The dynamical system (1) actually describes the mechanism that individuals tend to switch to stra that are doing well.
5.

ithe mentinditheth.0 1.0 eiry5i 8.821 -1.634 Td [(p)]TJ/F67 5.9776 Tf 3.986 -1.105 Td [(i)]TJ/F28 7.97F2

Continuous Strategy Space

1. Consider now games with pure strategies belonging to a continuum. For instance, this could be the aspiration level of a player or the size of an investment in economics (Hofbauer & Sigmund (2003)) or it might arise in situations where the pure strategies correspond to geographical points as in economic geography (Krugman (1996)).
2. On the other hand, in biology a continuous strategy space might correspond to some continuously varying trait such as the sex ratio in a birth or the virulence of an infection (Haccou & Iwasa (1998)).
3. Consider the case the strategy set is an arbitrary, not necessarily bounded, Borel set of $\mathbb{R}^N$, $N \geq 2$; hence strategies can be identified by $x \in \mathbb{R}^N$.
4. For the case of symmetric two-player games, the pay-off can be given by a Borel measurable function $f : \mathbb{R}^N \times \mathbb{R}^N \to \mathbb{R}$, where $f(x, y)$ is the pay-off for player 1 when she follows strategy x and player 2 plays strategy y.
5. A population is now characterized by its state, a probability measure P in the measure space $(\mathbb{R}^N, \mathcal{A})$ where $\mathcal{A}$ is the Borel algebra of subsets of $\mathbb{R}^N$. The average (mean) pay-off of a sub-population in state P against the overall population in state Q is given by the form

$$E(P, Q) := \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} f(x, y)\, Q(dy)\, P(dx).$$

Gaussian-type pay-off function

1.

The non-local model

For a bounded and smooth domain R^N then via integration by parts the non-local integro-differential dynamics equation (3) is approximated by the degenerate non-local parabolic equation

$$\frac{@u}{@t} = u \quad u + \int \quad jr \ uj^2 dx \ ; \quad x 2 \quad ; t > 0; \quad \int \quad u(x;t) = 1; \qquad (4)$$

(Kravvaritis, Papanicolaou, Xepapadeas & Yannacopoulos (2010)).

The non-local equation (4) is associated with initial condition

$$u(x;0) \int \quad 0 \quad u($$

Main estimates for the regularized problem (10)

Proposition 1 (Kowalczyk & Winkler (2017))

For all sufficiently small $\varepsilon \in (\varepsilon_j)_{j \in \mathbb{N}}$, problem (10) has a unique classical global-in-time solution $u_\varepsilon \in C^{2,1}(\overline{\Omega} \times [0,1))$.

The following lemma essentially derives a uniform pointwise bound for u_ε from a space-time integral estimate for $|\nabla u_\varepsilon|^2$:

Classi cation of initial data

Theorem 2 (Kaushansky & Wirdhan (2017))

Let u_0 satisfy (H1)-(H3), and let u and T_{max} denote the corresponding locally positive weak solution of (7)-(9)$_R$ as well as its maximal time of existence.
(i) If

Classi cation of initial data

Theorem 2 (Kaltenbacher & Winkler (2017))

Let u_0 satisfy (H1)-(H3), and let u and T_{max} denote the corresponding locally positive weak solution of (7)-(9)$_R$ as well as its maximal time of existence.
(i) If

Classi cation of initial data

Theorem 2 (Kaltenbacher & Winkler (2017))

Let u_0 satisfy (H1)-(H3), and let u and T_{max} denote the corresponding locally positive weak solution of (7)-(9)$_R$ as well as its maximal time of existence.
(i) If

Classi cation of initial data

Theorem 2 (Kaltenbacher & Winkler (2017))

Let u_0 satisfy (H1)-(H3), and let u and T_{max} denote the corresponding locally positive weak solution of (7)-(9)$_R$ as well as its maximal time of existence.
(i) If